



This is “Appendix B: Solubility-Product Constants (K_{sp}) for Compounds at 25°C”, appendix 2 from the book [Principles of General Chemistry \(index.html\)](#) (v. 1.0M).

This book is licensed under a [Creative Commons by-nc-sa 3.0](http://creativecommons.org/licenses/by-nc-sa/3.0/) license. See the license for more details, but that basically means you can share this book as long as you credit the author (but see below), don't make money from it, and do make it available to everyone else under the same terms.

This content was accessible as of December 29, 2012, and it was downloaded then by [Andy Schmitz](#) (<http://lardbucket.org>) in an effort to preserve the availability of this book.

Normally, the author and publisher would be credited here. However, the publisher has asked for the customary Creative Commons attribution to the original publisher, authors, title, and book URI to be removed. Additionally, per the publisher's request, their name has been removed in some passages. More information is available on this project's [attribution page](http://2012books.lardbucket.org/attribution.html?utm_source=header).

For more information on the source of this book, or why it is available for free, please see [the project's home page](#) (<http://2012books.lardbucket.org/>). You can browse or download additional books there.

Chapter 26

Appendix B: Solubility-Product Constants (K_{sp}) for Compounds at 25°C

Compound Name	Compound Formula	K_{sp}
Aluminum phosphate	AlPO_4	9.84×10^{-21}
Barium bromate	$\text{Ba}(\text{BrO}_3)_2$	2.43×10^{-4}
Barium carbonate	BaCO_3	2.58×10^{-9}
Barium chromate	BaCrO_4	1.17×10^{-10}
Barium fluoride	BaF_2	1.84×10^{-7}
Barium iodate	$\text{Ba}(\text{IO}_3)_2$	4.01×10^{-9}
Barium nitrate	$\text{Ba}(\text{NO}_3)_2$	4.64×10^{-3}
Barium sulfate	BaSO_4	1.08×10^{-10}
Barium sulfite	BaSO_3	5.0×10^{-10}
Beryllium hydroxide	$\text{Be}(\text{OH})_2$	6.92×10^{-22}
Bismuth arsenate	BiAsO_4	4.43×10^{-10}
Bismuth iodide	BiI_3	7.71×10^{-19}
Cadmium carbonate	CdCO_3	1.0×10^{-12}
Cadmium fluoride	CdF_2	6.44×10^{-3}
Cadmium hydroxide	$\text{Cd}(\text{OH})_2$	7.2×10^{-15}
Cadmium iodate	$\text{Cd}(\text{IO}_3)_2$	2.5×10^{-8}
Cadmium phosphate	$\text{Cd}_3(\text{PO}_4)_2$	2.53×10^{-33}
Cadmium sulfide	CdS	8.0×10^{-27}
Calcium carbonate	CaCO_3	3.36×10^{-9}
Calcium fluoride	CaF_2	3.45×10^{-11}
Calcium hydroxide	$\text{Ca}(\text{OH})_2$	5.02×10^{-6}
Calcium iodate	$\text{Ca}(\text{IO}_3)_2$	6.47×10^{-6}

Chapter 26 Appendix B: Solubility-Product Constants (K_{sp}) for Compounds at 25°C

Compound Name	Compound Formula	K_{sp}
Calcium phosphate	$\text{Ca}_3(\text{PO}_4)_2$	2.07×10^{-33}
Calcium sulfate	CaSO_4	4.93×10^{-5}
Cesium perchlorate	CsClO_4	3.95×10^{-3}
Cesium periodate	CsIO_4	5.16×10^{-6}
Cobalt(II) arsenate	$\text{Co}_3(\text{AsO}_4)_2$	6.80×10^{-29}
Cobalt(II) hydroxide	$\text{Co}(\text{OH})_2$	5.92×10^{-15}
Cobalt(II) phosphate	$\text{Co}_3(\text{PO}_4)_2$	2.05×10^{-35}
Copper(I) bromide	CuBr	6.27×10^{-9}
Copper(I) chloride	CuCl	1.72×10^{-7}
Copper(I) cyanide	CuCN	3.47×10^{-20}
Copper(I) iodide	CuI	1.27×10^{-12}
Copper(I) thiocyanate	CuSCN	1.77×10^{-13}
Copper(II) arsenate	$\text{Cu}_3(\text{AsO}_4)_2$	7.95×10^{-36}
Copper(II) oxalate	CuC_2O_4	4.43×10^{-10}
Copper(II) phosphate	$\text{Cu}_3(\text{PO}_4)_2$	1.40×10^{-37}
Copper(II) sulfide	CuS	6.3×10^{-36}
Europium(III) hydroxide	$\text{Eu}(\text{OH})_3$	9.38×10^{-27}
Gallium(III) hydroxide	$\text{Ga}(\text{OH})_3$	7.28×10^{-36}
Iron(II) carbonate	FeCO_3	3.13×10^{-11}
Iron(II) fluoride	FeF_2	2.36×10^{-6}
Iron(II) hydroxide	$\text{Fe}(\text{OH})_2$	4.87×10^{-17}
Iron(III) hydroxide	$\text{Fe}(\text{OH})_3$	2.79×10^{-39}
Iron(III) sulfide	FeS	6.3×10^{-18}
Lanthanum iodate	$\text{La}(\text{IO}_3)_3$	7.50×10^{-12}
Lead(II) bromide	PbBr_2	6.60×10^{-6}
Lead(II) carbonate	PbCO_3	7.40×10^{-14}
Lead(II) chloride	PbCl_2	1.70×10^{-5}
Lead(II) fluoride	PbF_2	3.3×10^{-8}

Chapter 26 Appendix B: Solubility-Product Constants (K_{sp}) for Compounds at 25°C

Compound Name	Compound Formula	K_{sp}
Lead(II) hydroxide	$Pb(OH)_2$	1.43×10^{-20}
Lead(II) iodate	$Pb(IO_3)_2$	3.69×10^{-13}
Lead(II) iodide	PbI_2	9.8×10^{-9}
Lead(II)selenite	$PbSeO_4$	1.37×10^{-7}
Lead(II) sulfate	$PbSO_4$	2.53×10^{-8}
Lead(II) sulfide	PbS	8.0×10^{-28}
Lithium carbonate	Li_2CO_3	8.15×10^{-4}
Lithium fluoride	LiF	1.84×10^{-3}
Lithium phosphate	Li_3PO_4	2.37×10^{-11}
Magnesium carbonate	$MgCO_3$	6.82×10^{-6}
Magnesium fluoride	MgF_2	5.16×10^{-11}
Magnesium hydroxide	$Mg(OH)_2$	5.61×10^{-12}
Magnesium phosphate	$Mg_3(PO_4)_2$	1.04×10^{-24}
Manganese(II) carbonate	$MnCO_3$	2.24×10^{-11}
Manganese(II) iodate	$Mn(IO_3)_2$	4.37×10^{-7}
Mercury(I) bromide	Hg_2Br_2	6.40×10^{-23}
Mercury(I) carbonate	Hg_2CO_3	3.6×10^{-17}
Mercury(I) chloride	Hg_2Cl_2	1.43×10^{-18}
Mercury(I) fluoride	Hg_2F_2	3.10×10^{-6}
Mercury(I) iodide	Hg_2I_2	5.2×10^{-29}
Mercury(I) oxalate	$Hg_2C_2O_4$	1.75×10^{-13}
Mercury(I) sulfate	Hg_2SO_4	6.5×10^{-7}
Mercury(I) thiocyanate	$Hg_2(SCN)_2$	3.2×10^{-20}
Mercury(II) bromide	$HgBr_2$	6.2×10^{-20}
Mercury (II) iodide	HgI_2	2.9×10^{-29}
Mercury(II) sulfide (red)	HgS	4×10^{-53}
Mercury(II) sulfide (black)	HgS	1.6×10^{-52}
Neodymium carbonate	$Nd_2(CO_3)_3$	1.08×10^{-33}

Chapter 26 Appendix B: Solubility-Product Constants (K_{sp}) for Compounds at 25°C

Compound Name	Compound Formula	K_{sp}
Nickel(II) carbonate	NiCO_3	1.42×10^{-7}
Nickel(II) hydroxide	Ni(OH)_2	5.48×10^{-16}
Nickel(II) iodate	$\text{Ni(IO}_3)_2$	4.71×10^{-5}
Nickel(II) phosphate	$\text{Ni}_3(\text{PO}_4)_2$	4.74×10^{-32}
Palladium(II) thiocyanate	Pd(SCN)_2	4.39×10^{-23}
Potassium hexachloroplatinate	K_2PtCl_6	7.48×10^{-6}
Potassium perchlorate	KClO_4	1.05×10^{-2}
Potassium periodate	KIO_4	3.71×10^{-4}
Praseodymium hydroxide	Pr(OH)_3	3.39×10^{-24}
Rubidium perchlorate	RbClO_4	3.00×10^{-3}
Scandium fluoride	ScF_3	5.81×10^{-24}
Scandium hydroxide	Sc(OH)_3	2.22×10^{-31}
Silver(I) acetate	AgCH_3CO_2	1.94×10^{-3}
Silver(I) arsenate	Ag_3AsO_4	1.03×10^{-22}
Silver(I) bromate	AgBrO_3	5.38×10^{-5}
Silver(I) bromide	AgBr	5.35×10^{-13}
Silver(I) carbonate	Ag_2CO_3	8.46×10^{-12}
Silver(I) chloride	AgCl	1.77×10^{-10}
Silver(I) chromate	Ag_2CrO_4	1.12×10^{-12}
Silver(I) cyanide	AgCN	5.97×10^{-17}
Silver(I) iodate	AgIO_3	3.17×10^{-8}
Silver(I) iodide	AgI	8.52×10^{-17}
Silver(I) oxalate	$\text{Ag}_2\text{C}_2\text{O}_4$	5.40×10^{-12}
Silver(I) phosphate	Ag_3PO_4	8.89×10^{-17}
Silver(I) sulfate	Ag_2SO_4	1.20×10^{-5}
Silver(I) sulfide	Ag_2S	6.3×10^{-50}
Silver(I) sulfite	Ag_2SO_3	1.50×10^{-14}
Silver(I) thiocyanate	AgSCN	1.03×10^{-12}

Chapter 26 Appendix B: Solubility-Product Constants (K_{sp}) for Compounds at 25°C

Compound Name	Compound Formula	K_{sp}
Strontium arsenate	$Sr_3(AsO_4)_2$	4.29×10^{-19}
Strontium carbonate	$SrCO_3$	5.60×10^{-10}
Strontium fluoride	SrF_2	4.33×10^{-9}
Strontium iodate	$Sr(IO_3)_2$	1.14×10^{-7}
Strontium sulfate	$SrSO_4$	3.44×10^{-7}
Thallium(I) bromate	$TlBrO_3$	1.10×10^{-4}
Thallium(I) bromide	$TlBr$	3.71×10^{-6}
Thallium(I) chloride	$TlCl$	1.86×10^{-4}
Thallium(I) chromate	Tl_2CrO_4	8.67×10^{-13}
Thallium(I) iodate	$TlIO_3$	3.12×10^{-6}
Thallium(I) iodide	TlI	5.54×10^{-8}
Thallium(I) thiocyanate	$TlSCN$	1.57×10^{-4}
Thallium(III) hydroxide	$Tl(OH)_3$	1.68×10^{-44}
Tin(II) hydroxide	$Sn(OH)_2$	5.45×10^{-27}
Tin(II) sulfide	SnS	1.0×10^{-25}
Yttrium carbonate	$Y_2(CO_3)_3$	1.03×10^{-31}
Yttrium fluoride	YF_3	8.62×10^{-21}
Yttrium hydroxide	$Y(OH)_3$	1.00×10^{-22}
Yttrium iodate	$Y(IO_3)_3$	1.12×10^{-10}
Zinc arsenate	$Zn_3(AsO_4)_2$	2.8×10^{-28}
Zinc carbonate	$ZnCO_3$	1.46×10^{-10}
Zinc fluoride	ZnF_2	3.04×10^{-2}
Zinc hydroxide	$Zn(OH)_2$	3×10^{-17}
Zinc selenide	$ZnSe$	3.6×10^{-26}
Zinc sulfide (wurtzite)	ZnS	1.6×10^{-24}
Zinc sulfide (sphalerite)	ZnS	2.5×10^{-22}

Chapter 26 Appendix B: Solubility-Product Constants (K_{sp}) for Compounds at 25°C

Source of data: *CRC Handbook of Chemistry and Physics*, 84th Edition (2004); sulfide data from *Lange's Handbook of Chemistry*, 15th Edition (1999).